

Package leaflet: Information for the patient

# COVERSYL<sup>®</sup> 4mg Tablets

Perindopril tert-butylamine

کوریسل<sup>®</sup> ۴ میلی گرام ٹیبلٹس  
بیرینڈوپریل ٲرٹری بوتائل امین

**Read all of this leaflet carefully before you start taking this medicine because it contains important information for you.**

- Keep this leaflet. You may need to read it again.
  - If you have any further questions, ask your doctor, pharmacist, or nurse.
  - This medicine has been prescribed for you only. Do not pass it on to others. It may harm them, even if their signs of illness are the same as yours.
- If you get any of the side effects, talk to your doctor, or pharmacist or nurse. This includes any possible side effects not listed in this leaflet. See section 4.

#### What is in this leaflet:

1. What Coversyl 4mg is and what it is used for
2. What you need to know before you take Coversyl 4mg
3. How to take Coversyl 4mg
4. Possible side effects
5. How to store Coversyl 4mg
6. Content of the pack and other information

#### 1. What Coversyl 4mg is and what it is used for

Coversyl 4mg is an angiotensin converting enzyme (ACE) inhibitor. These work by widening the blood vessels, which makes it easier for your heart to pump blood through them.

#### Coversyl 4mg is used:

- to treat **high blood pressure** (hypertension),
- to treat **heart failure** (a condition where the heart is unable to pump enough blood to meet the body's needs),
- to reduce the risk of cardiac events, such as heart attack, in patients with **stable coronary artery disease** (a condition where the blood supply to the heart is reduced or blocked) and who have already had a heart attack and/or an operation to improve the blood supply to the heart by widening the vessels that supply it.

#### 2. What you need to know before you take Coversyl 4mg

##### Do not take Coversyl 4mg

- if you are allergic to perindopril, or any of the other ingredients of this medicine (listed in Section 6), or to any other ACE inhibitor.
- if you have experienced symptoms such as wheezing, swelling of the face, tongue or throat, intense itching or severe skin rashes with previous ACE inhibitor treatment or if you or a member of your family have had these symptoms in any other circumstances (a condition called angioedema).
- if you are more than 3 months pregnant (It is also better to avoid Coversyl 4mg in early pregnancy – see pregnancy section).
- if you have diabetes or impaired kidney function and you are treated with a blood pressure lowering medicine containing aiskiren.
- if you are having dialysis or any other type of blood filtration. Depending on the machine that is used, Coversyl 4mg may not be suitable for you,
- if you have kidney problems where the blood supply to your kidneys is reduced (renal artery stenosis).
- if you have taken or are currently taking sacubitril/valsartan, a medicine for heart failure, as the risk of angioedema (rapid swelling under the skin in an area such as the throat) is increased (see "Warning and Precaution" and "Other medicines and Coversyl 4mg").

##### Warning and precaution

If any of the following apply to you please talk to your doctor, pharmacist or nurse before taking Coversyl 4mg if you:

- have aortic stenosis (narrowing of the main blood vessel leading from the heart) or hypertrophic cardiomyopathy (heart muscle disease) or renal artery stenosis (narrowing of the artery supplying the kidney with blood),
- have any other heart problems,
- have liver problems,
- have kidney problems or if you are receiving dialysis,
- have abnormally increased levels of a hormone called aldosterone in your blood (primary aldosteronism),
- suffer from a collagen vascular disease such as systemic lupus erythematosus or scleroderma,
- have diabetes,
- are on a salt restricted diet or use salt substitutes which contain potassium,
- are to undergo anaesthesia and/or major surgery,
- are to undergo LDL apheresis (which is removal of cholesterol from your blood by a machine),
- are going to have desensitisation treatment to reduce the effects of an allergy to bee or wasp stings,
- have recently suffered from diarrhoea or vomiting, or are dehydrated,
- have been told by your doctor that you have an intolerance to some sugars,
- are taking any of the following medicines used to treat high blood pressure:
  - an "angiotensin II receptor blocker" (ARBs) (also known as sartans - for example valsartan, telmisartan, irbesartan), in particular if you have diabetes-related kidney problems.
  - aiskiren.

Your doctor may check your kidney function, blood pressure, and the amount of electrolytes (e.g. potassium) in your blood at regular intervals.

See also information under the heading "Do not take Coversyl 4mg".

- are of black origin since you may have a higher risk of angioedema and this medicine may be less effective in lowering your blood pressure than in non-black patients.
- are taking any of the following medicines, the risk of angioedema is increased:

- racecadotril (used to treat diarrhea)
- sirolimus, everolimus, temsirolimus and other drugs belonging to the class of so-called mTor inhibitors (used to avoid rejection of transplanted organs and for cancer)
- sacubitril (available as fixed-dose combination with valsartan), used to treat long-term heart failure
- linagliptin, saxagliptin, sitagliptin, vildagliptin and other drugs belonging to the class of the also called gliptins (used to treat diabetes).

##### Angioedema

Angioedema (a severe allergic reaction with swelling of the face, lips, tongue or throat with difficulty in swallowing or breathing) has been reported in patients treated with ACE inhibitors, including Coversyl 4mg. This may occur at any time during treatment. If you develop such symptoms, you should stop taking Coversyl 4mg and see a doctor immediately. See also section 4.

You must tell your doctor if you think that you are (or might become) pregnant. Coversyl 4mg is not recommended in early pregnancy, and must not be taken if you are more than 3 months pregnant, as it may cause serious harm to your baby if used at that stage (see pregnancy section).

##### Children and adolescent

The use of perindopril in children and adolescents up to the age of 18 years is not recommended.

##### Other medicines and Coversyl 4mg

Please tell your doctor or pharmacist if you are taking, have recently taken or might take any other medicines.

Treatment with Coversyl 4mg can be affected by other medicines. Your doctor may need to change your dose and/or to take other precautions. These include:

- other medicines for high blood pressure, including angiotensin II receptor blocker (ARB), aiskiren (see also information under the headings "Do not take Coversyl 4mg" and "Warning and precaution"), or diuretics (medicines which increase the amount of urine produced by the kidneys),
- potassium-sparing drugs (e.g. triamterene, amiloride), potassium supplements or potassium-containing salt substitutes, other drugs which can increase potassium in your body (such as heparin, a medicine used to thin blood to prevent clots; trimethoprim and co-trimoxazole also known as trimethoprim/sulfamethoxazole for infections caused by bacteria),
- potassium-sparing drugs used in the treatment of heart failure: eplerenone and spironolactone at doses between 12,5 mg to 50 mg by day,
- lithium for mania or depression,
- non-steroidal anti-inflammatory drugs (e.g. ibuprofen) for pain relief or high dose acetylsalicylic acid, a substance presents in many medicines used to relieve pain and lower fever, as well as to prevent blood clotting,
- medicines to treat diabetes (such as insulin or metformin),
- baclofen (used to treat muscle stiffness in diseases such as multiple sclerosis),
- medicines to treat mental disorders such as depression, anxiety, schizophrenia etc (e.g. tricyclic antidepressants, antipsychotics),
- immunosuppressants (medicines which reduce the defence mechanism of the body) used for the treatment of auto-immune disorders or following transplant surgery (e.g. ciclosporin, tacrolimus),
- trimethoprim (for the treatment of infections),
- estramustine (used in cancer therapy),
- medicines, which are most often used to treat diarrhea (racecadotril) or avoid rejection of transplanted organs (sirolimus, everolimus, temsirolimus and other drugs belonging to the class of so-called mTor inhibitors). See section "Warnings and precautions",
- sacubitril/valsartan (used to treat long-term heart failure). See sections "Do not take Coversyl 4mg" and "Warnings and precautions".
- allopurinol (for the treatment of gout),
- procainamide (for the treatment of an irregular heart beat),
- vasodilators including nitrates (products that make the blood vessels become wider),
- medicines used for the treatment of low blood pressure, shock or asthma (e.g. ephedrine, noradrenaline or adrenaline),
- gold salts, especially with intravenous administration (used to treat symptoms of rheumatoid arthritis).

##### Coversyl 4mg with food and drink

It is preferable to take Coversyl 4mg before a meal.

##### Pregnancy and breast-feeding

If you are pregnant or breast-feeding, think you may be pregnant or are planning to have a baby, ask your doctor or pharmacist for advice before taking this medicine.

##### Pregnancy

You must tell your doctor if you think that you are (or might become) pregnant. Your doctor will normally advise you to stop taking Coversyl 4mg before you become pregnant or as soon as you know you are pregnant and will advise you to take another medicine instead of Coversyl 4mg. Coversyl 4mg is not recommended in early pregnancy, and must not be taken when more than 3 months pregnant, as it may cause serious harm to your baby if used after the third month of pregnancy.

##### Breastfeeding

Tell your doctor if you are breast-feeding or about to start breast-feeding. Coversyl 4mg is not recommended for mothers who are breast-feeding,

## Package leaflet: Information for the patient

and your doctor may choose another treatment for you if you wish to breast-feed, especially if your baby is newborn, or was born prematurely.

**Driving and using machines**

Coversyl 4mg usually does not affect alertness but dizziness or weakness due to low blood pressure may occur in certain patients. If you are affected in this way, your ability to drive or to operate machinery may be impaired.

**Coversyl 4mg contains lactose**

If you have been told by your doctor that you have an intolerance to some sugars, contact your doctor before taking this medicinal product.

**3. How to take Coversyl 4mg**

Always take this medicine exactly as your doctor or pharmacist has told you. Check with your doctor or pharmacist if you are not sure.

Swallow your tablet with a glass of water, preferably at the same time each day, in the morning, before a meal. Your doctor will decide on the correct dose for you.

The recommended dosages are as follows:

**High blood pressure:** the usual starting and maintenance dose is 4 mg once daily. After one month, this can be increased to 8 mg once a day if required. 8 mg a day is the maximum recommended dose for high blood pressure.

If you are 65 or older, the usual starting dose is 2 mg once a day. After a month this can be increased to 4 mg once a day and then if necessary to 8 mg once daily.

**Heart failure:** the usual starting dose is 2 mg once daily. After two weeks, this can be increased to 4 mg once a day, which is the maximum recommended dose for heart failure.

**Stable coronary artery disease:** the usual starting dose is 4 mg once daily. After two weeks, this can be increased to 8 mg once daily, which is the maximum recommended dose in this indication.

If you are 65 or older, the usual starting dose is 2 mg once a day. After a week this can be increased to 4 mg once a day and after a further week to 8 mg once daily.

**Use in children and adolescent**

Use in children and adolescent is not recommended.

**If you take more Coversyl 4mg than you should**

If you take too many tablets, contact your nearest accident and emergency department or tell your doctor immediately. The most likely effect in case of overdose is low blood pressure which can make you feel dizzy or faint. If this happens, lying down with the legs raised can help.

**If you forget to take Coversyl 4mg**

It is important to take your medicine every day as regular treatment works better. However, if you forget to take a dose of Coversyl 4mg, take the next dose at the usual time. Do not take a double dose to make up for a forgotten dose.

**If you stop taking Coversyl 4mg**

As the treatment with Coversyl 4mg is usually life-long, you should discuss with your doctor before stopping this medicinal product.

If you have any further questions on the use of this product, ask your doctor or pharmacist.

**4. Possible side effects**

Like all medicines, this medicine can cause side effects, although not everybody gets them.

**Stop taking the medicinal product and see a doctor immediately, if you experience any of the following side effects that can be serious:**

- swelling of the face, lips, mouth, tongue or throat, difficulty in breathing (angioedema) (See section 2 "Warning and precaution") (Uncommon - may affect up to 1 in 100 people),
- severe dizziness or fainting due to low blood pressure (Common - may affect up to 1 in 10 people),
- unusual fast or irregular heartbeat, chest pain (angina) or heart attack (Very rare - may affect up to 1 in 10,000 people),
- weakness of arms or legs, or problems speaking which could be a sign of a possible stroke (Very rare - may affect up to 1 in 10,000 people),
- sudden wheeziness, chest pain, shortness of breath, or difficulty in breathing (bronchospasm) (Uncommon - may affect up to 1 in 100 people),
- nflamed pancreas which may cause severe abdominal and back pain accompanied with feeling very unwell (Very rare - may affect up to 1 in 10,000 people),
- yellowing of the skin or eyes (jaundice) which could be a sign of hepatitis (Very rare - may affect up to 1 in 10,000 people),
- skin rash which often starts with red itchy patches on your face, arms or legs (erythema multiforme) (Very rare - may affect up to 1 in 10,000 people).

**Tell your doctor if you notice any of the following side effects:**

Common (may affect up to 1 in 10 people):

- headache,
- dizziness,
- vertigo,
- pins and needles,
- vision disturbances,
- tinnitus (sensation of noises in the ears),
- cough,
- shortness of breath (dyspnoea),
- gastro-intestinal disorders (nausea, vomiting, abdominal pain, taste disturbances, dyspepsia (indigestion), diarrhoea, constipation),
- allergic reactions (such as skin rashes, itching),
- muscle cramps,
- feeling of tiredness.

Uncommon (may affect to 1 in 100 people):

- depression,
- mood swings,

- sleep disturbances,
- dry mouth,
- intense itching or severe skin rashes,
- formation of blister clusters over the skin,
- kidney problems,
- impotence,
- sweating,
- excess of eosinophils (a type of white blood cells),
- somnolence,
- fainting,
- palpitations,
- tachycardia,
- vasculitis (inflammation of blood vessels),
- photosensitivity reaction (increased sensitivity of the skin to sun),
- arthralgia (joint pain),
- myalgia (muscle pain),
- chest pain,
- malaise,
- oedema peripheral,
- fever,
- fall,
- change in laboratory parameters: high blood level of potassium reversible on discontinuation, low level of sodium, hypoglycaemia (very low blood sugar level) in case of diabetic patients, increase blood urea, and increased blood creatinine.

Rare (may affect up to 1 in 1000 people):

- acute renal failure,
- dark urine, feeling sick (nausea) or being sick (vomiting), muscle cramps, confusion and seizures. These may be symptoms of a condition called SIADH (inappropriate antidiuretic hormone secretion).
- decreased or absent urine output,
- flushing,
- psoriasis worsening,
- changes in laboratory parameters: Increased level of liver enzymes, high level of serum bilirubin.

Very rare (may affect up to 1 in 10,000 people):

- confusion,
- eosinophilic pneumonia (a rare type of pneumonia),
- rhinitis (blocked up or runny nose),
- changes in blood values such as a lower number of white and red blood cells, lower haemoglobin, lower number of blood platelets.

If you have these symptoms contact your doctor as soon as possible.

Frequency not known (cannot be estimated from available data):  
Discoloration, numbness and pain in fingers or toes (Raynaud's phenomenon).

**Reporting of side effects**

If you get any side effects, talk to your doctor or pharmacist or nurse. This includes any possible side effects not listed in this leaflet.

You can also report side effects directly to company website [www.servier.com.pk](http://www.servier.com.pk) or to Drug Regulatory Authority of Pakistan [www.dra.gov.pk](http://www.dra.gov.pk). By reporting side effects you can help provide more information on the safety of this medicine.

**5. How to store Coversyl 4mg**

Keep this medicine out of the sight and reach of children.

Do not use this medicine after the expiry date which is stated on the carton and blister. The expiry date refers to the last day of that month. Do not store above 30°C.

Do not throw away any medicine via wastewater or household waste. Ask your pharmacist how to throw away of medicines you no longer used. These measures will help to protect the environment.

**6. Content of the pack and other information****What Coversyl 4mg contains**

- The active substance is perindopril tert-butylamine. One tablet contains 3.338 mg perindopril (corresponding to 4 mg perindopril tert-butylamine).
- The other ingredients are: microcrystalline cellulose, lactose monohydrate, hydrophobic colloidal silica, magnesium stearate.

**What Coversyl 4mg looks like and contents of the pack**

Coversyl 4mg tablets are Light Green, rod-shaped, scored on both faces. The tablet can be divided into equal halves.

The tablets are available in blister packs (PVC/Aluminium) of 10 tablets.

**Marketing Authorisation Holder and Manufacturer**

Servier Research & Pharmaceuticals [Pakistan] (Pvt) Ltd  
9-KM Sheikhpura Road Lahore Pakistan

**MARKETING AUTHORISATION NUMBER**

011458

**DATE OF FIRST AUTHORISATION**

31-07-1990

**DATE OF REVISION OF THE TEXT**

February 2022

Variation Reference No: MU 41121

04058  
Version-06



**Manufactured by:**  
**Servier Research & Pharmaceuticals [Pakistan] (Pvt.) Ltd.**  
9-Km Sheikhpura Road, Lahore - Pakistan  
Under Licence **Les Laboratoires Servier - France**